

Teacher Preparation Notes for “Were the babies switched? – The Genetics of Blood Types”¹

In this minds-on, hands-on activity, students use simple chemicals to simulate blood type tests and carry out genetic analyses to determine whether hospital staff accidentally switched two babies born on the same day. Students learn about the genetics of ABO blood types, multiple alleles of a single gene, and codominance. Next, students analyze the genetics of skin color in order to understand how fraternal twins can have different skin colors. In this analysis, students learn about incomplete dominance and how a single phenotypic characteristic can be influenced by multiple genes and the environment.²

As background for this activity, students should have a basic understanding of:

- dominant and recessive alleles (heterozygous individuals have the same phenotype as homozygous dominant individuals)
- how Punnett squares summarize inheritance.

To provide this background you may want to use the first four pages of “Introduction to Genetics – Similarities and Differences between Family Members”

(<https://serendipstudio.org/exchange/bioactivities/geneticsFR>) or the first 2.5 pages of "Genetics" (https://serendipstudio.org/sci_edu/waldron/#genetics).

Table of Contents

Learning Goals – pages 1-2

Supplies, Suggestions for Implementation, and Preparation – pages 2-5

General Instructional Suggestions – page 5

Biology Background and Suggestions for Discussion

The Genetics of Blood Types – pages 5-9

Analysis of the Genetics of Skin Color – pages 9-11

Additional Genetics Learning Activities – page 11

Optional Immunobiology Addition to Student Handout – pages 12-13

Learning Goals Related to National Standards

In accord with the Next Generation Science Standards³ and A Framework for K-12 Science Education⁴:

- Students will gain understanding of two Disciplinary Core Ideas:
 - LS1.A: Structure and Function – "All cells contain genetic information in the form of DNA molecules. Genes are regions in the DNA that contain the instructions that code for the formation of proteins."
 - LS3.A: Inheritance of Traits – "The instructions for forming species' characteristics are carried in DNA."
- Students will engage in two Scientific Practices:
 - Constructing Explanations. “Apply scientific ideas, principles, and/or evidence to provide an explanation of phenomena...”
 - Analyzing And Interpreting Data. “Analyze data using tools, technologies, and/or models (e.g. computational, mathematical) in order to make valid and reliable scientific claims...”

¹ By Drs. Jennifer Doherty and Ingrid Waldron, Department of Biology, University of Pennsylvania, 2022. These Teacher Preparation Notes and the Student Handout are available at <https://serendipstudio.org/exchange/waldron/bloodtests>.

² The same genetic concepts are covered in an analysis and discussion activity (“Were the babies switched?” in “Soap Opera Genetics – Genetics to Resolve Family Arguments”; <https://serendipstudio.org/exchange/bioactivities/SoapOperaGenetics>).

³ <https://www.nextgenscience.org/sites/default/files/HS%20LS%20topics%20combined%206.13.13.pdf>

⁴ http://www.nap.edu/catalog.php?record_id=13165

- This activity provides the opportunity to discuss the Crosscutting Concept, Structure and Function. "Cause and effect relationships can be suggested and predicted for complex natural and human designed systems by examining what is known about smaller scale mechanisms within the system."
- This activity helps to prepare students for the Performance Expectation, HS-LS3-1, "Ask questions to clarify relationships about the role of DNA and chromosomes in coding the instructions for characteristic traits passed from parents to offspring."

Additional Content Learning Goals

- Each person has one of the four blood types: A, B, AB, and O. These blood types refer to the presence or absence of two different versions of a carbohydrate molecule (A and B) on the surface of red blood cells.
- Genes code for proteins which influence a person's characteristics. The ABO blood type gene codes for a protein enzyme that can attach a carbohydrate molecule to the surface of red blood cells. This gene has three alleles: the E^A allele codes for a version of the enzyme that attaches the type A carbohydrate; the E^B allele codes for a version of the enzyme that attaches the type B carbohydrate; and the e allele codes for an inactive protein that does not attach either carbohydrate.
- Each person inherits one allele of this gene from his/her mother and a second allele from his/her father. In the cells that are precursors of red blood cells, both inherited alleles provide the instructions for making proteins.
- In a person who has the $E^A E^B$ genotype, both alleles are transcribed, so the cells produce both the version of the enzyme that puts type A carbohydrates on the surface of red blood cells and the version of the enzyme that puts type B carbohydrates on the surface of red blood cells. Therefore, the person has type AB blood. This is an example of codominance, in which two alleles of a gene each have a different observable effect on the phenotype of the heterozygous individual.
- In a heterozygous person with the $E^A e$ or $E^B e$ genotype, the single copy of the E^A or E^B allele in each cell codes for enough enzyme to result in type A or type B blood, respectively. Thus, the e allele is recessive relative to the E^A or E^B alleles.
- In incomplete dominance, the phenotype of a heterozygous individual is intermediate between the phenotypes of the two different types of homozygous individuals (observed for quantitative traits such as skin color).
- Many characteristics are influenced by multiple genes and environmental factors.

Supplies, Suggestions for Implementation, and Preparation

Note: Throughout this section we will refer to the seven people listed in the table on the next page as the subjects.

Supplies

(See subsequent pages for information about the needed amounts and types of the first three items.)

- Synthetic blood
- Solutions with synthetic anti-A antibodies and anti-B antibodies
- Small dropper bottles (can be reused in multiple classes)
- Non-porous testing surfaces suitable for mixing samples of blood with antibody solution, e.g. blood-typing trays, microscope slides or white or transparent plastic lids (can be washed and reused in multiple classes); each student group will need a testing surface large enough for 14 tests (two tests for each subject) or several smaller testing surfaces.

- One marker for each student group to identify the 14 specific spots to test for the type A antigen and for the type B antigen for each subject.
- Toothpicks for mixing blood and antibody solution (Each student group will need 14 toothpicks, or they can use both ends of 7 toothpicks.)
- Containers such as plastic cups or water bottles to use as trash containers so the students can throw away their toothpicks immediately after use to avoid contamination

To determine the required amount of synthetic blood and antibodies, you will need to decide what blood types you will assign to each subject. You may want to vary the blood types for each subject in different classes, in order to maintain some variety and suspense. Each column of this table presents a possible combination of blood types for each subject.

Examples of Blood Type Combinations You Can Use

	1	2	3	4	5	6	7	8	9	10
Danielle (mother of twins)	AB	AB	AB	AB	O	A	B	AB	A	A
Michael (father of twins)	O	A	B	AB	AB	AB	AB	AB	O	A
Denise (mother of daughter)	A	A	A	A	B	B	B	B	A	A
Earnest (father of daughter)	B	B	B	B	A	A	A	A	B	B
Michael Jr. (boy twin)	A	A	A	A	B	B	B	B	A	A
Baby girl 1 (girl twin, according to hospital)	B	B	B	B	A	A	A	A	O	O
Baby girl 2 (daughter of Denise and Earnest, according to hospital)	O	O	O	O	O	O	O	O	B	B

You can make other combinations, provided that:

- Michael Jr. can be the son of Danielle and Michael
- One of the baby girls could be a daughter of one of the couples and could not be a daughter of the other couple. The other baby girl could be a daughter of the other couple.

Option 1. Colored milk as the simulated blood and water or vinegar as the simulated antibody solutions:

You can make simulated blood by combining 25 mL of milk with red food coloring until the solution is bright red (about 15 drops), and then adding a drop of green food coloring for a dark red color.⁵ You will need different anti-A and anti-B simulated solutions for each subject, depending on what blood type the sample is supposed to be.

Type of Blood	Simulated anti-A solution contains:	Simulated anti-B solution contains:
A	White vinegar	Water
B	Water	White vinegar
AB	White vinegar	White vinegar
O	Water	Water

⁵ We have had success with fat-free milk. We have not tried other types of milk, but they would probably work fine.

We recommend that you set up seven stations, one for each of the seven subjects, with the blood sample and the anti-A and anti-B solutions. Before class you should prepare seven bottles with the simulated blood, labeled with the subject's name or Baby girl 1 or 2. You will also need the corresponding bottles of simulated anti-A and anti-B solutions. We recommend that you label each bottle of anti-A and anti-B solution with the name of the subject which will help you keep track of which pair of antibody solutions goes with each subject.

You can use the various blood type combinations shown on the previous page. You can estimate amounts of simulated blood and antibody solution needed using the following information:

- For the blood type test for each person, you will need 6 drops of blood, 3 drops of anti-A antibody solution, and 3 drops of anti-B antibody solution.
- There are approximately 15-20 drops in each milliliter of solution.

You will want extra of each solution, in order to be prepared for student error such as using too many drops of a solution.

Option 2. Commercial simulated blood and antibody solutions:

To determine the amount of synthetic blood, antibody solutions and dropper bottles you will need, you should choose one of these recommendations for implementation (or decide on your own approach).

- Give each student group at their lab table:
 - 7 bottles with the blood samples, each labeled with the name of one of the subjects or Baby girl 1 or 2
 - 1 bottle with the anti-A antibody solution and another bottle with the anti-B antibody solution (each labeled appropriately)
- Or you can set up three stations:
 - one where each student group will get the blood samples for each of the seven subjects
 - one where they will get the anti-A antibody solution
 - one where they will get the anti-B antibody solution.

Please note that the procedure on page 3 of the Student Handout is written for the colored milk simulated blood. If you are purchasing simulated blood, you may need to modify the procedure; for example, you can use only two drops each of simulated blood and simulated antibody solution.

You can purchase kits (and/or refills) from:

- Carolina (http://www.carolina.com/blood-typing/carolina-abo-rh-blood-typing-with-synthetic-blood-kit/FAM_700101.pr?question=700101) (We recommend you not use the Rh antiserum included in this kit.)
- Ward's Science <https://www.wardsci.com/store/product/10424340/simulated-abo-blood-typing-kit>⁶

These kits have additional supplies such as some dropper bottles and testing trays. You may want to contact these companies to verify that their kits have the blood types and quantities you will need. This table shows the amounts of antibody solution and blood type solution you will need per student group for each of the blood type combinations in the first table on the previous page .

⁶ We recommend against purchasing these supplies from www.Neoscience.com. One teacher has reported problems with the Ward's kit. Please send feedback about the Carolina or Ward's kits to iwaldron@upenn.edu .

**Approximate Amount (mL) Needed of Each Type of Solution
for Each Student Group for Blood Type Combinations Listed on page 3**

	1	2	3	4	5	6	7	8	9	10
Anti-A Antibody Solution	1	1	1	1	1	1	1	1	1	1
Anti-B Antibody Solution	1	1	1	1	1	1	1	1	1	1
A Blood	0.6	0.8	0.6	0.6	0.6	0.8	0.6	0.6	0.8	1.1
B Blood	0.6	0.6	0.8	0.6	0.6	0.6	0.8	0.6	0.6	0.6
AB Blood	0.3	0.3	0.3	0.6	0.3	0.3	0.3	0.6	0.0	0.0
O Blood	0.6	0.3	0.3	0.3	0.6	0.3	0.3	0.3	0.6	0.3

You will want more of each solution, in order to be prepared for student error such as using too many drops of a solution.

General Instructional Suggestions

To maximize student learning, we recommend that you have your students complete groups of related questions in the Student Handout in pairs and then have a class discussion of student answers to each group of related questions. In each discussion, you can probe student thinking and help them to develop a sound understanding of the concepts and information covered before moving on to the next part of the activity.

The PDF of the Student Handout shows the correct format; please check this if you use the Word document to make revisions.

If you would like to have a key, please send your request to iwaldron@upenn.edu. The following paragraphs provide additional instructional suggestions and background information – some for inclusion in your class discussions and some to provide you with relevant background that may be useful for your understanding and/or for responding to student questions.


Biology Background and Suggestions for Discussion

In the introductory section of the Student Handout, Danielle insists on blood type tests for the parents and babies to determine whether the baby girls were switched. Modern methods use DNA testing to determine biological relatedness; these results are much more definitive than testing blood types (http://en.wikipedia.org/wiki/Parental_testing).

The Genetics of Blood Types

Question 1 provides the opportunity to reinforce student understanding that genes code for proteins which influence an organism’s characteristics. Thus, the three alleles of the gene for ABO blood types provide the instructions for making three versions of the protein enzyme that attaches the carbohydrate molecules to lipid or protein molecules in the plasma membrane of red blood cells (see figure below).⁷


⁷ There is more than one version of each of these three alleles, but this is not mentioned in the Student Handout.



(Modified from https://www.researchgate.net/figure/The-structure-of-terminal-sugars-in-the-A-B-and-O-blood-antigens-Gal-Galactose_fig7_4074726)

For the ABO blood type gene:

- The E^A allele codes for the version of the protein enzyme that attaches the type A carbohydrate to protein or lipid molecules in the plasma membrane that surrounds each red blood cell. The resulting glycoproteins and glycolipids have the carbohydrates on the outer surface of the cell membrane (see figure below).⁸
- The E^B allele codes for a different version of this enzyme that attaches the type B carbohydrate to membrane proteins and lipids.
- The e allele codes for an inactive version of this protein enzyme.



(https://cnx.org/resources/3a229fd6909a56722463c77805188a3f/Figure_05_01_01.jpg)

The function of the blood type carbohydrate molecules is unknown. In general, people who have Type O blood with neither Type A nor Type B carbohydrates are as healthy as people who have

⁸ The E and e in the symbols for the allele stands for enzyme. Usually, these are designated I or i , which stands for isoagglutinin; isoagglutinin refers to an antigen that can stimulate the production of antibodies in other members of the same species; these antibodies result in agglutination (clumping) upon exposure to the antigen. The type A and type B glycoproteins and glycolipids are often referred to as type A and type B antigens, because they can trigger an immune response.

the Type A and/or Type B carbohydrates. Different blood types are correlated with certain illnesses and vary in frequency in different ethnic groups, but the reasons are unknown (<https://www.webmd.com/a-to-z-guides/ss/slideshow-how-your-blood-type-affects-your-health>).

Before question 2, you may want to ask your students these two questions.

- “Explain why a person with the **ee** genotype has blood type **O**.”
- “What is a dominant allele?”

For a dominant-recessive pair of alleles, heterozygous individuals have the same phenotype as individuals who are homozygous for the dominant allele. You may also want to point out that recessive alleles often code for a nonfunctional protein. In a heterozygous individual, a single dominant allele can code for enough functional protein to result in the same phenotype as the phenotype of the homozygous dominant individual. For example, the **e** allele codes for a nonfunctional protein, and the **e** allele is recessive relative to the **E^A** or **E^B** alleles because, in a heterozygous individual, the single dominant **E^A** or **E^B** allele codes for enough functional enzyme to result in the same blood type as observed in a homozygous dominant individual.

This activity helps students to understand the molecular basis for codominance. Each cell in the body contains two copies of each gene and typically both alleles are transcribed. Thus, at the molecular level, the alleles of most genes are codominant.⁹ For example, the **E^AE^B** genotype results in the production of both the version of the enzyme that puts type A carbohydrate on red blood cells and the version of the enzyme that puts type B carbohydrate on red blood cells. Therefore, the **E^AE^B** genotype results in type AB blood.


As you discuss student answers to questions 5-6 you may want to reinforce student understanding that:

- meiosis and fertilization result in new combinations of alleles, so children may have different blood types (and other phenotypic characteristics) than their parents have
- transmission of genes via meiosis and fertilization result in similarities between offspring and parents.

Testing Blood Types, Procedure, Results and Interpretation


The type A and type B glycoproteins and glycolipids can stimulate the body to produce an immune response, including antibodies. Normally, your body does *not* make antibodies against any molecules that are part of your body. This is useful because antibodies against antigens that are part of your body can trigger harmful reactions. As would be expected, a person does not make antibodies against the blood type carbohydrates present on their red blood cells. However, a person with type A blood does make anti-B antibodies. The figure below shows the type of antibodies in the blood plasma for each blood type. These antibodies are believed to be induced by ubiquitous environmental antigens such as antigens on influenza virus or gut bacteria. (An antigen is a molecule that can stimulate the production of antibodies.)

⁹ For example, a person who is heterozygous for the allele for normal hemoglobin and sickle cell hemoglobin has both types of hemoglobin in their red blood cells. Due to the normal hemoglobin in the red blood cells of a heterozygous person, the hemoglobin molecules very rarely clump into rods that distort the shape of the red blood cells, so the heterozygous person very rarely develops the symptoms of sickle cell anemia. The sickle cell hemoglobin in the red blood cells of a heterozygous person inhibits the reproduction of the malaria parasite in the red blood cells, so the heterozygous person is protected against severe malaria infections.



(https://biology-forums.com/gallery/14755_01_10_12_7_59_21_9228594.jpeg)

The figure below shows what would happen if a person with type A blood were given a transfusion of type B blood. The anti-B antibodies in the recipient's blood would cause the donated type B red blood cells to clump together and burst. This could block blood vessels and could result in kidney failure. This type of transfusion reaction can be fatal. Notice that only people with type AB blood are universal recipients (i.e., can safely receive transfusions of type A, B, AB or O blood); this illustrates codominance at the phenotypic level.



This figure shows what would happen if a person with type A blood were given a transfusion of type B blood.
(https://biology-forums.com/gallery/14755_01_10_12_7_59_21_9228594.jpeg)


The relatively small quantity of anti-A or anti-B antibodies in a blood transfusion do not cause significant health problems. Therefore, type O blood can be safely given to people with type A, B, AB or O blood, since type O red blood cells do not have either A or B antigens. Also, people with type AB blood can safely receive transfusions of type A, B, AB or O blood, since people with type AB blood do not have anti-A or anti-B antibodies in their blood. The unique ability of people with type AB blood to safely receive transfusions of all four types of blood illustrates codominance at the phenotypic level.

To prevent a transfusion reaction, medical personnel test whether a person's blood is compatible with the donated blood before they give a transfusion. The ABO blood types are the major determinant of which type of blood will cause a transfusion reaction. However, determination of blood type is more complex than the ABO blood types discussed in this activity. For additional information on blood types, see:

- <http://www.ncbi.nlm.nih.gov/books/NBK2264/>.
- [https://bio.libretexts.org/Bookshelves/Human_Biology/Book%3A_Human_Biology_\(Wakim_and_Grewal\)/17%3A_Cardiovascular_System/17.6%3A_Blood_Types](https://bio.libretexts.org/Bookshelves/Human_Biology/Book%3A_Human_Biology_(Wakim_and_Grewal)/17%3A_Cardiovascular_System/17.6%3A_Blood_Types)
- <https://www.britannica.com/science/ABO-blood-group-system>

If you want your students to have a more extended introduction to the relevant immunobiology, you can replace the section, “Testing Blood Types”, on the top of page 3 of the Student Handout with the section, “Using Immune Responses to Test Blood Types”, provided on pages 12-13 of these Teacher Preparation Notes.

If you do this, you may want to mention that antibodies help to protect our bodies against infection by helping to destroy bacteria and viruses. For example, some types of antibodies bind to the antigens on the surface of bacteria that have infected a person’s body. The other end of each antibody can bind to a protein on the surface of a type of cell called a macrophage. This facilitates phagocytosis of the bacteria by the macrophage. The macrophage kills and digests the ingested bacteria.



For suggestions about implementation of the experimental procedure, see “Supplies, Suggestions for Implementation, and Preparation” on pages 2-5 of these Teacher Preparation Notes. The “Results” and “Interpretation” sections are straightforward if students have understood the previous sections and are familiar with Punnett squares.

Why do the twins look so different?

The bottom of page 4 of the Student Handout introduces the important distinction between identical (monozygotic) twins and fraternal (dizygotic) twins.

The top of page 5 of the Student Handout introduces a very simplified model of the genetic determination of skin color, with a single gene that illustrates incomplete dominance.¹⁰ The table below may be helpful for your class discussion of student answers to question 12.

Type of Dominance	Phenotype of Heterozygous Individual
Dominant-recessive pair of alleles	Same as phenotype of individual who is homozygous for the dominant allele
Codominance	Shows different observable phenotypic effects of both alleles; phenotype different from either homozygous individual
Incomplete dominance	Intermediate between phenotypes of the two types of homozygous individual (typically observed for quantitative traits); phenotype different from either homozygous individual

The bottom of page 5 of the Student Handout develops a more realistic and complex model of the genetic factors that influence skin color. During your discussion of student answers to question 14, you may want to explain that the genotype/phenotype table at the top of the page provides a very simplified introduction to the genetics of skin color. A person with a **Tt** genotype


¹⁰ Incomplete dominance can occur when each wild type allele produces a set dose of protein product and the phenotype is proportionate to the amount of protein. The Student Handout uses a capital letter and lowercase letter to indicate the two alleles for a gene with incomplete dominance; you may prefer to use an alternate notation such as **t/t⁺** or **T/T'**.

could have lighter or darker skin, depending on whether he or she:

- has alleles for other genes that contribute to lighter or darker skin color
- has developed a tan due to sun exposure or tanning booth use.

Similarly, two people with a **TT** genotype (or a **tt** genotype) could have different skin colors, depending on the alleles of the other genes that influence skin color and the amount of UV exposure. You will probably want to explain that scientists often begin with a simple model of a phenomenon, which is superseded by a more complex model as the scientists learn more.

You may want to further explain that, compared to the Punnett square in answer to question 13, this figure provides a more accurate representation of the Punnett square for inheritance of skin color. Even this relatively complex Punnett square is a simplified representation of reality, since it assumes a simple additive model with only two alleles for each gene and incomplete dominance for all of the alleles.



One gene that influences skin color codes for the enzyme tyrosinase, which is a crucial enzyme involved in the synthesis of melanin, the primary pigment in skin and hair. The normal allele of this gene codes for functional tyrosinase; other alleles code for defective, non-functional versions of this enzyme which result in albinism. The alleles for albinism are recessive because, even when there is only one copy of the normal allele, this allele codes for enough functioning enzyme to produce enough melanin to result in normal skin and hair color.

Another important gene that influences skin color is the MC1R gene which codes for the melanocortin receptor; when alpha melanocyte stimulating hormone binds to normal melanocortin receptor this stimulates melanocytes to produce melanin. More than 80 alleles of the MC1R gene have been identified, resulting in various levels of function of the melanocortin receptor and corresponding variation in skin tones. Heterozygotes for these alleles have intermediate skin color, between the lighter and darker homozygotes (called incomplete dominance or a dosage effect). The multiple alleles and the effects of incomplete dominance result in multiple different phenotypes for skin color (and hair color). Additional information on this gene is available at <https://ghr.nlm.nih.gov/gene/MC1R>.

Additional information on the complex genetics and molecular biology involved in regulation of skin color is available at <http://www.jbc.org/content/282/38/27557.full> and <http://hmg.oxfordjournals.org/content/18/R1/R9.full>.

This analysis provides the opportunity to reinforce student understanding that, often, an individual phenotypic characteristic is influenced by multiple alleles of multiple genes, as well as environmental factors. Our introductory genetics teaching often focuses on inheritance and phenotypic effects of single genes. However, this is only a beginning for understanding the genetics of most traits.

Additional Genetics Learning Activities

"Genetics – Major Concepts and Learning Activities"

(<https://serendipstudio.org/exchange/bioactivities/GeneticsConcepts>)

Part I summarizes key concepts in genetics. Part II presents common misconceptions. Part III recommends an integrated sequence of learning activities on the biological basis of genetics, plus seven human genetics learning activities. These learning activities develop student understanding of key concepts and counteract common misconceptions. Each of these recommended learning activities supports the Next Generation Science Standards ([NGSS](#)).¹¹


¹¹ <http://www.nextgenscience.org/>

Optional Immunobiology Addition to Student Handout

If you want a more extended introduction to the relevant immunobiology, you can insert this section in place of the shorter section, “Testing Blood Types”, on the top of page 3 of the Student Handout.

Using Immune Responses to Test Blood Types

The type A and type B carbohydrate molecules on the surface of red blood cells are called **antigens** because they can stimulate the body to produce (generate) **antibodies**. Each specific type of antibody binds to a specific antigen. For example, anti-B antibodies in the blood bind to type B antigens, but not to type A antigens.



Antibodies are molecules that help the immune system to identify and destroy invaders such as bacteria or viruses that have entered your body. When an antibody binds with an antigen on the surface of a cell, this typically results in damage to the cell. Fortunately, your body generally does not make antibodies against any antigens that are part of your body.

7. Fill in the blanks in this chart.

<p>If you have <u>type A</u> blood, you have:</p> <ul style="list-style-type: none"> • type A antigens on the surface of your red blood cells and • _____ antibodies in your blood. 	<p>If you have <u>type B</u> blood, you have:</p> <ul style="list-style-type: none"> • type B antigens on the surface of your red blood cells and • _____ antibodies in your blood. 	<p>If you have <u>type AB</u> blood, you have:</p> <ul style="list-style-type: none"> • both type ___ and type ___ antigens on the surface of your red blood cells and • neither anti-A nor anti-B antibodies in your blood. 	<p>If you have <u>type O</u> blood, you have:</p> <ul style="list-style-type: none"> • neither type of antigen on the surface of your red blood cells and • both _____ and _____ antibodies in your blood.

8. If a person with type A blood is given a transfusion of type B blood, the _____ antibodies in his/her blood will react with the type B antigens on the donated red blood cells and cause them to clump together and block blood vessels (see top figure on this page).

This type of transfusion reaction can be fatal. To prevent a transfusion reaction, doctors test whether donated blood is compatible with a person’s blood before they give a blood transfusion.

9. To test the blood type of each parent and each baby, you will:

- mix one sample of their blood with anti-A antibodies to test whether their blood has the type ____ antigen
- mix a second sample of their blood with anti-B antibodies to test whether their blood has the type ____ antigen.

If the sample has antigens that match the antibodies, this will result in clumping.

10. To prepare to interpret the blood type tests, fill in this chart.

Blood type	Will this blood type clump if mixed with	
	anti-A antibody?	anti-B antibody?
A		
B		
AB		
O		